

REVIEW ARTICLE

CURRENT CONCEPTS

Rhabdomyolysis and Acute Kidney Injury

Xavier Bosch, M.D., Ph.D., Esteban Poch, M.D., Ph.D.,
and Josep M. Grau, M.D., Ph.D.

From the Muscle Research Unit, Department of Internal Medicine (X.B., J.M.G.), and the Department of Nephrology (E.P.), Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, Barcelona. Address reprint requests to Dr. Grau at the Muscle Research Unit, Department of Internal Medicine, Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, Villarroel 170 08036-Barcelona, Spain, or at jmgrau@clinic.ub.es, or to Dr. Poch at the Department of Nephrology, Hospital Clínic, Villarroel 170, 08036-Barcelona, Spain, or at epoch@clinic.ub.es.

N Engl J Med 2009;361:62-72.
Copyright © 2009 Massachusetts Medical Society.

RHABDOMYOLYSIS — LITERALLY, THE DISSOLUTION OF STRIPED (SKELETAL) muscle — is characterized by the leakage of muscle-cell contents, including electrolytes, myoglobin, and other sarcoplasmic proteins (e.g., creatine kinase, aldolase, lactate dehydrogenase, alanine aminotransferase, and aspartate aminotransferase) into the circulation. Massive necrosis, which is manifested as limb weakness, myalgia, swelling, and, commonly, gross pigmenturia without hematuria, is the common denominator of both traumatic and nontraumatic rhabdomyolysis.^{1,2} Acute kidney injury is a potential complication of severe rhabdomyolysis, regardless of whether the rhabdomyolysis is the result of trauma or some other cause, and the prognosis is substantially worse if renal failure develops. In contrast, in less severe forms of rhabdomyolysis or in cases of chronic or intermittent muscle destruction — a condition sometimes called hyperCKemia — patients usually present with few symptoms and no renal failure. We review the pathophysiological characteristics and management of acute kidney injury associated with rhabdomyolysis.

There are eight commonly reported categories of rhabdomyolysis (Table 1). Exogenous agents that can be toxic to muscles, especially alcohol, illicit drugs, and lipid-lowering agents, are common nontraumatic causes. Recurrent episodes of rhabdomyolysis are often a sign of an underlying defect in muscle metabolism.^{1,3,4}

Acute rhabdomyolysis occasionally develops in patients with structural myopathies when they are performing strenuous exercise, are under anesthesia, have taken drugs that are toxic to muscles, or have viral infections.¹ When a diagnosis of acute rhabdomyolysis is suspected, histochemical, immunohistochemical, and mitochondrial respiration studies performed on a muscle-biopsy specimen may yield a specific diagnosis. It is important to wait several weeks or months after the clinical event to perform a biopsy, because the results of a biopsy will typically be uninformative at an early stage. Thus, the specimen may appear normal or show no specific findings other than necrosis during and early after the acute episode of rhabdomyolysis (Fig. 1).^{2,5}

The mechanisms involved in the pathogenesis of rhabdomyolysis are direct sarcolemmic injury (e.g., trauma) or depletion of ATP within the myocyte, leading to an unregulated increase in intracellular calcium.^{6,7} Sarcoplasmic calcium is strictly regulated by a series of pumps, channels, and exchangers that maintain low levels of calcium when the muscle is at rest and allow the increase that is necessary for actin–myosin binding and muscle contraction. Depletion of ATP impairs the function of these pumps, resulting in a persistent increase in sarcoplasmic calcium that leads to persistent contraction and energy depletion and the activation of calcium-dependent neutral proteases and phospholipases; the result is the eventual destruction of myofibrillar, cytoskeletal, and membrane proteins, followed by lysosomal digestion of fiber contents. Ultimately, the myofibrillar network breaks down, resulting in disintegration of the myocyte.² In the case of patients with rhabdomy-

Table 1. Major Categories and Commonly Reported Causes of Rhabdomyolysis.

Category	Commonly Reported Cause
Trauma	Crush syndrome
Exertion	Strenuous exercise, seizures, alcohol withdrawal syndrome
Muscle hypoxia	Limb compression by head or torso during prolonged immobilization or loss of consciousness,* major artery occlusion
Genetic defects	Disorders of glycolysis or glycogenolysis, including myophosphorylase (glycogenosis type V), phosphofructokinase (glycogenosis type VII), phosphorylase kinase (glycogenosis type VIII), phosphoglycerate kinase (glycogenosis type IX), phosphoglycerate mutase (glycogenosis type X), lactate dehydrogenase (glycogenosis type XI) Disorders of lipid metabolism, including carnitine palmitoyl transferase II, long-chain acyl-CoA dehydrogenase, short-chain L-3-hydroxyacyl-CoA dehydrogenase, medium-chain acyl-CoA dehydrogenase, very-long-chain acyl-CoA dehydrogenase, medium-chain 3-ketoacyl-CoA, thiolase† Mitochondrial disorders, including succinate dehydrogenase, cytochrome <i>c</i> oxidase, coenzyme Q10 Pentose phosphate pathway: glucose-6-phosphate dehydrogenase Purine nucleotide cycle: myoadenylate deaminase
Infections‡	Influenza A and B, coxsackievirus, Epstein–Barr virus, primary human immunodeficiency virus, legionella species <i>Streptococcus pyogenes</i> , <i>Staphylococcus aureus</i> (pyomyositis), clostridium
Body-temperature changes	Heat stroke, malignant hyperthermia, malignant neuroleptic syndrome, hypothermia
Metabolic and electrolyte disorders	Hypokalemia, hypophosphatemia, hypocalcemia, nonketotic hyperosmotic conditions, diabetic ketoacidosis
Drugs and toxins	Lipid-lowering drugs (fibrates, statins), alcohol, heroin, cocaine
Idiopathic (sometimes recurrent)	

* Rhabdomyolysis from this cause is associated with a crush syndrome–like mechanism.

† CoA denotes coenzyme A.

‡ In most cases, the mechanism is unclear.


olysis caused by trauma, additional injury results from ischemia reperfusion and inflammation by neutrophils that infiltrate damaged muscle.⁸

EPIDEMIOLOGY OF MYOGLOBINURIA-INDUCED ACUTE KIDNEY INJURY

Acute kidney injury associated with myoglobinuria is the most serious complication of both traumatic and nontraumatic rhabdomyolysis, and it may be life-threatening. Acute kidney injury as a complication of rhabdomyolysis is quite common, representing about 7 to 10% of all cases of acute kidney injury in the United States.^{4,9} The true incidence of acute kidney injury in rhabdomyolysis is difficult to establish owing to varying definitions and clinical scenarios. The reported incidence ranges from 13% to approximately 50%.^{9–11} In a study by Melli et al. involving 475 hospitalized patients with rhabdomyolysis, the incidence of acute kidney injury was 46%.¹⁰ Although rhabdomyolysis from any cause can lead to acute kidney injury, in this study, the incidence of acute kidney injury was higher among persons who used illicit drugs or abused alcohol and among

persons who had undergone trauma than among persons with muscle disease, and the incidence was particularly high among persons with more than one recognized causal factor.¹⁰

The outcome of rhabdomyolysis is usually good provided that there is no renal failure. Nevertheless, mortality data vary widely according to the study population and setting and the number and severity of coexisting conditions. In a study in which the incidence of vasculopathy leading to rhabdomyolysis as a result of limb ischemia was high, the overall mortality was 32%.¹² In contrast, the study by Melli et al. of hospitalized patients, in whom the abuse of illicit drugs and alcohol was the most frequently identified cause of rhabdomyolysis, showed a mortality of 3.4% among patients with acute kidney injury.¹⁰ Among patients in the intensive care unit, the mortality has been reported to be 59% when acute kidney injury is present and 22% when it is not present.^{13,14} Long-term survival among patients with rhabdomyolysis and acute kidney injury is reported to be close to 80%, and the majority of patients with rhabdomyolysis-induced acute kidney injury recover renal function.¹⁴



PATHOGENESIS OF MYOGLOBIN-INDUCED ACUTE KIDNEY INJURY

Myoglobinuria occurs only in the context of rhabdomyolysis. Myoglobin is a dark red 17.8-kDa protein that is freely filtered by the glomerulus, enters the tubule epithelial cell through endocytosis, and is metabolized. It appears in the urine only when the renal threshold of 0.5 to 1.5 mg of myoglobin per deciliter is exceeded and is grossly visible as reddish-brown (“tea-colored”) urine when serum myoglobin levels reach 100 mg per deciliter¹⁵; therefore, not all cases of rhabdomyolysis are associated with myoglobinuria.

Although the exact mechanisms by which rhabdomyolysis impairs the glomerular filtration rate are unclear, experimental evidence suggests that intrarenal vasoconstriction, direct and is-

chemic tubule injury, and tubular obstruction all play a role (Fig. 2).¹⁶ Myoglobin becomes concentrated along the renal tubules, a process that is enhanced by volume depletion and renal vasoconstriction, and it precipitates when it interacts with the Tamm–Horsfall protein, a process favored by acidic urine.¹⁷ Tubule obstruction occurs principally at the level of the distal tubules, and direct tubule cytotoxicity occurs mainly in the proximal tubules.

Myoglobin seems to have no marked nephrotoxic effect in the tubules unless the urine is acidic. Myoglobin is a heme protein; it contains iron, as ferrous oxide (Fe^{2+}), which is necessary for the binding of molecular oxygen. However, molecular oxygen can promote the oxidation of Fe^{2+} to ferric oxide (Fe^{3+}), thus generating a hydroxyl radical. This oxidative potential is counter-

acted by effective intracellular antioxidant molecules. However, cellular release of myoglobin leads to uncontrolled leakage of reactive oxygen species, and free radicals cause cellular injury. It has been suggested that heme and free iron-driven hydroxyl radicals are critical mediators of tubule damage owing to the protective effects of deferoxamine (an iron chelator) and glutathione.¹⁸ More recently, it has been shown that myoglobin itself can exhibit peroxidase-like enzyme activity that leads to uncontrolled oxidation of biomolecules, lipid peroxidation, and the generation of isoprostanes.¹⁹

Renal vasoconstriction is a characteristic feature of rhabdomyolysis-induced acute kidney injury and is the result of various combinations of several mechanisms. First, intravascular volume depletion due to fluid sequestration within damaged muscle promotes homeostatic activation of the renin-angiotensin system, vasopressin, and the sympathetic nervous system. Second, experimental studies have shown that there are additional vascular mediators in the reduction of renal blood flow, including endothelin-1, thromboxane A₂, tumor necrosis factor α ; and F₂-isoprostanes^{9,20}; a deficit in the vasodilator nitric oxide, which can be attributed to the scavenging effect of myoglobin in the renal microcirculation, has also been shown to be a mediator in the reduction in renal blood flow.¹⁶ Collectively, these vascular mediators appear to be locally stimulated by oxidant injury and leukocyte-mediated inflammation as a result of the endothelial dysfunction that is common to other forms of acute kidney injury.²¹


RENAL MANIFESTATIONS OF RHABDOMYOLYSIS

Patients with acute rhabdomyolysis usually present with pigmented granular casts, reddish-brown urine supernatant, and markedly raised serum creatine kinase. There is no defined threshold value of serum creatine kinase above which the risk of acute kidney injury is markedly increased. A very weak correlation between the peak creatine kinase value and the incidence of acute kidney injury or peak serum creatinine has been reported.^{10,11,22} The risk of acute kidney injury in rhabdomyolysis is usually low when creatine kinase levels at admission are less than 15,000 to 20,000 U per liter.^{12,13,23} Although acute kidney injury may be associated with creatine kinase values as low

as 5000 U per liter, this usually occurs when co-existing conditions such as sepsis, dehydration, and acidosis are present.¹¹ For example, in patients with chronic myopathies such as muscular dystrophies and inflammatory myopathies, acute kidney injury seldom develops unless a superimposed event is present. Patients with these chronic myopathies, on the other hand, may have moderately raised concentrations of plasma myoglobin but not overt myoglobinuria.²⁴ Myoglobinuria can be inferred if urinary dipstick testing shows a positive result for blood when there are no red cells in the sediment. This false positive result for blood occurs because the dipstick test is unable to distinguish between myoglobin and hemoglobin. The test has a sensitivity of 80% for the detection of rhabdomyolysis.¹⁰ Other causes of pigmented urine should be taken into consideration (Table 2).²⁵ Myoglobin is the true pathogenic factor in rhabdomyolysis-induced acute kidney injury but is seldom measured directly in urine or plasma. Serum myoglobin levels peak well before serum creatine kinase levels, and serum myoglobin has a rapid and unpredictable metabolism, which functions partly through the kidney but mainly outside the kidney (probably through the liver or spleen).²⁶ Therefore, measurement of serum myoglobin has a low sensitivity for the diagnosis of rhabdomyolysis.²⁷

Acute kidney injury associated with rhabdomyolysis often leads to a more rapid increase in plasma creatinine than do other forms of acute kidney injury. However, this finding may reflect the overrepresentation of young, muscular men among patients with rhabdomyolysis rather than increased creatinine or creatine release from injured muscle.^{14,28,29} Similarly, a low ratio of blood urea nitrogen to creatinine is often seen in patients with rhabdomyolysis. Rhabdomyolysis-induced acute kidney injury frequently causes oliguria and occasionally causes anuria.

Another characteristic feature of rhabdomyolysis-induced acute kidney injury that is different from the manifestation of other forms of acute tubular necrosis is the frequent, but not universal, presence of a low fractional excretion of sodium (<1%), perhaps reflecting the primacy of preglomerular vasoconstriction and tubular occlusion rather than tubular necrosis.³⁰ The fractional excretion of sodium is a measurement of the percentage of filtered sodium that is excreted in the urine, and low levels in patients with acute kidney injury are an indication of the relative



integrity of tubular functions. However, when ischemic or toxic acute tubular necrosis is established, both urinary sodium and the fractional excretion of sodium are raised.

Electrolyte abnormalities that occur as a result of the release of cellular components often accompany and determine the severity of rhabdomyolysis-induced acute kidney injury. Because

Table 2. Causes and Microscopic Features of Red and Brown Urine.

Cause	Results of Test for Blood in Fresh Urine [*]	Sediment ^{†‡}	Supernatant [‡]
Hematuria	+ to ++++	Red	Yellow
Myoglobinuria	+ to ++++	Normal	Red to brown
Hemoglobinuria	+ to ++++	Normal	Red to brown
Porphyria	Negative	Normal	Red
Bile pigments	Negative	Normal	Brown
Food and drugs [§]	Negative	Normal	Red to brown

* Urine was tested with the use of a dipstick test. This is a semiquantitative test of the number of erythrocytes per microliter. Results range from + (5 to 10 erythrocytes per microliter) to ++++ (approximately 250 erythrocytes per microliter).

† Normal refers to white or yellow in color, unremarkable in the absence of cells, crystals, or cylinders.

‡ The sediment and supernatant were examined after centrifugation of 10 to 15 ml of urine at 1500 to 3000 rpm for 5 minutes.

§ Food and drugs that can cause red urine include beets, blackberries, rhubarb, food coloring, fava beans, phenolphthalein, rifampin, doxorubicin, deferoxamine, chloroquine, ibuprofen, and methyldopa. Those that can cause brown urine include levodopa, metronidazole, nitrofurantoin, iron sorbitol, chloroquine, and methyldopa.

they may precede the acute kidney injury, electrolyte levels should be measured as soon as rhabdomyolysis is diagnosed. The electrolyte abnormalities that can occur with rhabdomyolysis include hyperkalemia (which can be rapidly increasing), hyperphosphatemia, hyperuricemia, high anion-gap metabolic acidosis, and hypermagnesemia mainly when renal failure is present.^{4,15,22,31} High levels of phosphate can bind to calcium, and deposition of calcium-phosphate complexes in soft tissues can occur. In addition, hyperphosphatemia inhibits 1 α -hydroxylase, thus limiting the formation of calcitriol (1,25-dihydroxyvitamin D₃), the active form of vitamin D. Hyperkalemia is an early manifestation of rhab-

domyolysis, and serum potassium can occasionally reach life-threatening levels both in patients with severe traumatic rhabdomyolysis and in those with nontraumatic rhabdomyolysis.^{15,31} Hyperuricemia is also usually present owing to the liberation of nucleosides from injured muscle and can contribute to renal tubule obstruction since uric acid is insoluble and may precipitate in acidic urine.

Hypocalcemia is a common complication of rhabdomyolysis and usually results from calcium entering the ischemic and damaged muscle cells and from the precipitation of calcium phosphate with calcification in necrotic muscle. Hypercalcemia associated with recovery of renal function is unique to rhabdomyolysis-induced acute kidney injury and results from the mobilization of calcium that was previously deposited in muscle, the normalization of hyperphosphatemia, and an increase in calcitriol.³²

TREATMENT AND PREVENTION

Patients with rhabdomyolysis that is associated with acute kidney injury usually present with a clinical picture of volume depletion that is due to the sequestration of water in injured muscles. Therefore, the main step in managing the condition (Table 3) remains the early, aggressive repletion of fluids; patients often require about 10 liters of fluid per day,³¹ with the amount administered depending on the severity of the rhabdomyolysis.³³ There are no randomized trials that have evaluated fluid repletion in patients with the crush syndrome resulting from injuries sus-

Figure 2 (facing page). Pathophysiological Mechanisms in Rhabdomyolysis-Induced Acute Kidney Injury.

Fluid sequestration in injured muscle induces volume depletion and consequent activation of the sympathetic nervous system (SNS), antidiuretic hormone (ADH), and the renin-angiotensin system (RAS), all of which favor vasoconstriction and renal salt and water conservation. In addition, myoglobin-induced oxidative injury increases vasoconstrictors and decreases vasodilators. Kidney injury results from a combination of ischemia due to renal vasoconstriction, direct tubular toxicity mediated by myoglobin-associated oxidative injury (inset, lower right), tubular damage due to ischemia, and distal tubule obstruction due to precipitation of the Tamm-Horsfall protein-myoglobin complex (inset, lower left) in addition to sloughed tubular cells forming cellular cast. As in acute kidney injury due to other causes, endothelial dysfunction and local inflammation contribute to tissue damage and organ dysfunction. ET denotes endothelin, F₂ IP F₂ isoprostanes, NO nitric oxide, THP Tamm-Horsfall protein, TNF- α tumor necrosis factor α , TxA₂ thromboxane A₂, and VC vasoconstriction.

Table 3. Steps in the Prevention and Treatment of Rhabdomyolysis-Induced Acute Kidney Injury.

<p>Check for extracellular volume status, central venous pressure, and urine output.*</p> <p>Measure serum creatine kinase levels. Measurement of other muscle enzymes (myoglobin, aldolase, lactate dehydrogenase, alanine aminotransferase, and aspartate aminotransferase) adds little information relevant to the diagnosis or management.</p> <p>Measure levels of plasma and urine creatinine, potassium and sodium, blood urea nitrogen, total and ionized calcium, magnesium, phosphorus, and uric acid and albumin; evaluate acid–base status, blood-cell count, and coagulation.</p> <p>Perform a urine dipstick test and examine the urine sediment.</p> <p>Initiate volume repletion with normal saline promptly at a rate of approximately 400 ml per hour (200 to 1000 ml per hour depending on the setting and severity), with monitoring of the clinical course or of central venous pressure.</p> <p>Target urine output of approximately 3 ml per kilogram of body weight per hour (200 ml per hour).</p> <p>Check serum potassium level frequently.</p> <p>Correct hypocalcemia only if symptomatic (e.g., tetany or seizures) or if severe hyperkalemia occurs.</p> <p>Investigate the cause of rhabdomyolysis.</p> <p>Check urine pH. If it is less than 6.5, alternate each liter of normal saline with 1 liter of 5% dextrose or 0.45% saline plus 100 mmol of bicarbonate. Avoid potassium and lactate-containing solutions.</p> <p>Consider treatment with mannitol (up to 200 g per day and cumulative dose up to 800 g). Check for plasma osmolality and plasma osmolal gap. Discontinue if diuresis (>20 ml per hour) is not established.</p> <p>Maintain volume repletion until myoglobinuria is cleared (as evidenced by clear urine or a urine dipstick testing result that is negative for blood).</p> <p>Consider renal-replacement therapy if there is resistant hyperkalemia of more than 6.5 mmol per liter that is symptomatic (as assessed by electrocardiography), rapidly rising serum potassium, oliguria (<0.5 ml of urine per kilogram per hour for 12 hours), anuria, volume overload, or resistant metabolic acidosis (pH <7.1).</p>
--

* In the case of the crush syndrome (e.g., earthquake, building collapse), institute aggressive volume repletion promptly before evacuating the patient.

tained in a disaster such as an earthquake. However, most, if not all, reports show that patients in whom acute kidney injury developed had a longer delay in receiving supportive therapy than did patients in whom acute kidney injury did not develop (Table 4).^{8,23,31,33,39} Therefore, early, aggressive volume repletion is crucial in patients with the crush syndrome.^{34,35}

Although the need for volume repletion is established, the composition of the fluid used for repletion remains controversial. Some investigators recommend administering sodium bicarbonate, which results in an alkaline urine, as first proposed by Bywaters and Beall,^{8,39,40} whereas others argue against this approach and favor normal or 0.45% saline solution.¹⁵ The three empirical advantages of alkalization that have been noted are based on studies in animal models of rhabdomyolysis. First, it is known that precipitation of the Tamm–Horsfall protein–myoglobin complex is increased in acidic urine.¹⁷ Second, alkalization inhibits reduction–oxidation (redox) cycling of myoglobin and lipid peroxidation in rhabdomyolysis, thus ameliorating tubule injury.⁴¹ Third, it has been shown that metmyoglobin induces vasoconstriction only in an acidic

medium in the isolated perfused kidney.⁴² The principal, and probably the only, disadvantage of alkalization is the reduction in ionized calcium, which can exacerbate the symptoms of the initial hypocalcemic phase of rhabdomyolysis.

The clinical benefits of alkalization as compared with simple volume repletion are not firmly established. Comparative studies usually have small sample sizes and show a combination of measures (e.g., alkalization plus mannitol) that preclude an analysis of the effectiveness of the particular single measure^{34–38} (Table 4). In one study, renal outcomes did not differ significantly between patients treated with bicarbonate plus mannitol and those treated with saline alone, although peak serum creatine kinase values were below 5000 U per liter, a finding indicating that the degree of injury was mild, making treatment effect difficult to appreciate.³⁶ In the largest study of patients who had undergone trauma (2083 patients), rhabdomyolysis developed in 85% of the patients, and administration of bicarbonate plus mannitol did not prevent renal failure, the need for dialysis, or death in the sample as a whole, although the results suggested that it might be beneficial in patients with peak creatine kinase

Table 4. Comparative Studies on Preventive and Therapeutic Regimens in Rhabdomyolysis.

Study	Study Design	Patient Group	No. in Sample	Therapeutic Strategy	Outcome in Patients with Acute Kidney Injury
Shimazu et al. ³⁴	Retrospective	Patients with the crush syndrome	14	Late vs. early initiation of therapy; high (>10 liters for 48 hours) vs. low volume of hydration	Better if therapy initiated early; high volume of hydration better
Gunal et al. ³⁵	Retrospective	Patients with the crush syndrome	16	Early vs. late treatment with normal saline followed immediately by bicarbonate	Better if treatment initiated early
Homsi et al. ³⁶	Retrospective	Patients in the intensive care unit	24	Normal saline vs. normal saline plus bicarbonate and mannitol	No difference
Brown et al. ³⁷	Retrospective	Patients with trauma	2083	Normal saline vs. bicarbonate plus mannitol	No difference
Cho et al. ³⁸	Prospective, randomized	Patients with intoxication from doxylamine	28	Ringer's lactate vs. normal saline; bicarbonate if urine pH is <6.5	No effect on peak creatine kinase level or recovery with Ringer's lactate as compared with normal saline; more bicarbonate needed with normal saline than with Ringer's lactate

values of more than 30,000 U per liter.³⁷ In a randomized, prospective trial of fluid repletion with Ringer's lactate as compared with normal saline in patients with rhabdomyolysis attributed to doxylamine intoxication, 28 patients were randomly assigned to receive one of the solutions.³⁸ Sodium bicarbonate was added in both groups if the urine pH was less than 6.5 after 12 hours of aggressive volume repletion. Peak creatine kinase levels were less than 10,000 U per liter, and it appears that acute kidney injury did not develop in any of the patients, although these data were not reported. Whatever the real, consistent benefits of urine alkalinization in patients with rhabdomyolysis, there is evidence that massive infusion of normal saline alone can contribute to metabolic acidosis, mainly owing to the dilution of serum bicarbonate with a solution relatively high in chloride ions, generating hyperchloremic metabolic acidosis with observed reductions in serum pH of as much as 0.30 units.⁴³ Therefore, administration of both normal saline and sodium bicarbonate seems to be a reasonable approach when fluid is being replenished in patients with rhabdomyolysis, especially patients with metabolic acidosis (Table 3). If sodium bicarbonate is used, urine pH and serum bicarbonate, calcium, and potassium levels should be monitored, and if the urine pH does not rise after 4 to 6 hours of treatment or if symptomatic hypocalcemia develops, alkalinization should be discontinued and hydration continued with normal saline.

The use of diuretics remains controversial, but it is clear that it should be restricted to patients in whom the fluid repletion has been achieved. Mannitol may have several benefits: as an osmotic diuretic, it increases urinary flow and the flushing of nephrotoxic agents through the renal tubules; as an osmotic agent, it creates a gradient that extracts fluid that has accumulated in injured muscles and thus improves hypovolemia; finally, it is a free-radical scavenger.^{4,8,20} Most data on the action of mannitol come from studies in animals, which collectively show that the protective effect of mannitol may be attributable to its osmotic diuretic action rather than to the other mechanisms.⁴⁴ No randomized, controlled trial has supported the evidence-based use of mannitol, and some clinical studies suggest no beneficial effects.^{36,37} In addition, high accumulated doses of mannitol (>200 g per day or accumulated doses of >800 g) have been associated with acute kidney injury due to renal vasoconstriction and tubular toxicity, a condition known as osmotic nephrosis.^{45,46} However, many experts continue to suggest that mannitol should be used to prevent and treat rhabdomyolysis-induced acute kidney injury and relieve compartmental pressure.^{20,45-47} During the time mannitol is being administered, plasma osmolality and the osmolal gap (i.e., the difference between the measured and calculated serum osmolality) should be monitored frequently and therapy discontinued if adequate diuresis is not achieved or if the osmolal gap rises above 55 mOsm per kilogram.⁴⁶ Loop

Table 5. Approach to the Management of Hyperkalemia (Serum Potassium ≥ 5.5 mmol per Liter) in Rhabdomyolysis.

<p>Check for potassium levels every 4 hours in cases of severe rhabdomyolysis (creatinine kinase level $>60,000$ to $80,000$ U per liter) or suspected systemic toxin. Treat rapidly rising potassium levels aggressively.</p> <p>Obtain an electrocardiogram and check for severe manifestations (QRS interval widening, small P waves, severe arrhythmias thought to be caused by high levels of potassium). Consider cardiac monitoring and admission to an intensive care unit if the potassium level is higher than 6 mmol per liter, if there are abnormalities on the electrocardiogram, or if rhabdomyolysis is severe, with rapidly rising potassium.</p> <p>Check for plasma calcium levels. Hypocalcemia seriously aggravates the adverse electrical effects of hyperkalemia.</p> <p>If the electrocardiogram shows severe irregularities, administer calcium chloride or calcium gluconate by intravenous infusion. Consider slow continuous infusion if hypocalcemia is present. Anticipate possible hypercalcemia in late rhabdomyolysis. Do not mix with bicarbonate solutions.</p> <p>If potassium level is higher than 6 mmol per liter, shift potassium into cells. Serum potassium will be lowered approximately 10 to 30 minutes after the following measures are performed, and the effect will last for 2 to 6 hours.</p> <p>Administer insulin and glucose by means of a slow intravenous push; monitor glucose with the use of fingerstick testing.</p> <p>Administer a β_2-adrenergic agonist such as albuterol, 10 to 20 mg in 4 ml of normal saline by inhalation of aerosol over 10 minutes. Do not use as a single measure; combine with glucose and insulin for additive effect.</p> <p>Administer sodium bicarbonate if the patient has acidemia. This treatment may worsen the manifestations of hypocalcemia, and the efficacy is not as consistent as that with insulin and glucose or albuterol. Do not use as a single measure.</p> <p>Remove potassium from the body with the use of either resins or dialysis as indicated; the use of diuretics is optional.</p> <p>Administer cation-exchange resin (sodium polystyrene sulfonate) orally or as a retention enema (avoid sorbitol in such cases and avoid after surgery).</p> <p>Perform hemodialysis if the above measures fail or if severe renal failure or severe hyperkalemia develops. Consider hemodialysis when rhabdomyolysis is associated with marked tissue breakdown and rapidly rising serum potassium levels. Check serum potassium levels 4 hours after hemodialysis, since a rebound increase can occur. Previous measures of potassium shift into cells may decrease the efficiency of hemodialysis with respect to removal of potassium.</p> <p>Administer loop diuretics such as furosemide, but only after the patient's fluid level has been expanded.</p>
--

diuretics also increase urinary flow and may decrease the risk of myoglobin precipitation, but no study has shown a clear benefit in patients with rhabdomyolysis. Therefore, loop diuretics in rhabdomyolysis-induced acute kidney injury should be used in the same manner as that recommended in acute kidney injury that is due to other causes.^{47,48}

The electrolyte abnormalities associated with rhabdomyolysis-induced acute kidney injury must be treated promptly; the correction of hyperkalemia, which occurs very early in the course of the disease, is especially important (Table 5).⁴⁹ Agents that cause a shift of potassium from the extracellular to the intracellular space (e.g., hypertonic glucose and bicarbonate) are effective only temporarily, and the only means of removing potassium from the body is diuresis (effective kaliuresis), the use of intestinal potassium binders, or dialysis.^{4,8,9,15} In contrast, early hypocalcemia should not be treated unless it is symptomatic or unless severe hyperkalemia is present. Calcium-containing chelators should be used with caution to treat hyperphosphatemia, since the calcium load could increase the precipitation of calcium phosphate in injured muscle.^{4,8,9,15}

When acute kidney injury is severe enough to produce refractory hyperkalemia, acidosis, or vol-

ume overload, renal-replacement therapy is indicated, principally with intermittent hemodialysis, which can correct electrolyte abnormalities rapidly and efficiently.^{8,9,47} Conventional hemodialysis does not remove myoglobin effectively owing to the size of the protein and is therefore usually mandated by renal indications. However, owing to the pathogenic role of myoglobin in rhabdomyolysis-induced acute kidney injury, preventive extracorporeal elimination has been studied. Although plasmapheresis has been shown to have no effect on outcomes or on the myoglobin burden of the kidneys,⁵⁰ continuous venovenous hemofiltration or hemodiafiltration has shown some efficacy in removing myoglobin, principally with the use of super high-flux filters and high volumes of ultrafiltration (convection).⁵¹ However, the evidence is mainly from isolated case reports, and the effect on outcomes is unknown. In addition, some studies have shown that the half-life of serum myoglobin does not differ significantly between patients who are treated conservatively and those who receive continuous venovenous hemodiafiltration.²⁷ Until randomized studies are performed, preventive hemofiltration cannot be recommended.

The use of antioxidants and free-radical scav-

engers (e.g., pentoxifylline, vitamin E, and vitamin C) may be justified in the treatment or prevention of myoglobinuric acute kidney injury,^{8,52} as suggested by small case series, case reports, and various experimental studies of myoglobinuria, but controlled studies evaluating their efficacy are lacking.

Supported by grants from Fondo Investigaciones Sanitarias (FIS 05/0015, to Dr. Poch; and FIS 04/0464, to Dr. Grau), Instituto de Salud Carlos III, Red Renal de Investigación Cooperativa (ISCI-III-Retic-RD06, to Dr. Poch), Suport Grup de Recerca (05/300, to Dr. Grau), and the Center for Biomedical Research on Rare Diseases, Instituto de Salud Carlos III, Barcelona (to Dr. Grau).

No potential conflict of interest relevant to this article was reported.

We thank Assumpta Violan, M.D., for her invaluable technical assistance in preparing the original draft of Figure 2.

REFERENCES

- Tein I, DiMauro S, Rowland LP. Myoglobinuria. In: Rowland LP, DiMauro S, eds. *Myopathies. Handbook of clinical neurology*. Vol. 62. Amsterdam: Elsevier Science Publishers, 1992:553-93.
- Warren JD, Blumberg PC, Thompson PD. Rhabdomyolysis: a review. *Muscle Nerve* 2002;25:332-47.
- Allison RC, Bedsole DL. The other medical causes of rhabdomyolysis. *Am J Med Sci* 2003;326:79-88.
- Bagley WH, Yang H, Shah KH. Rhabdomyolysis. *Intern Emerg Med* 2007;2:210-8.
- Fernandez-Sola J, Grau JM, Pedro-Botet JC, et al. Nontraumatic rhabdomyolysis: a clinical and morphological analysis of 53 cases. *Med Clin (Barc)* 1988;90:199-202. (In Spanish.)
- Giannoglou GD, Chatzizisis YS, Misirli G. The syndrome of rhabdomyolysis: pathophysiology and diagnosis. *Eur J Intern Med* 2007;18:90-100.
- Wroegemann K, Pena SD. Mitochondrial calcium overload: a general mechanism for cell-necrosis in muscle diseases. *Lancet* 1976;1:672-4.
- Vanholder R, Sever MS, Ereik E, Lameire N. Rhabdomyolysis. *J Am Soc Nephrol* 2000;11:1553-61.
- Holt SG, Moore KP. Pathogenesis and treatment of renal dysfunction in rhabdomyolysis. *Intensive Care Med* 2001;27:803-11.
- Melli G, Chaudhry V, Cornblath DR. Rhabdomyolysis: an evaluation of 475 hospitalized patients. *Medicine (Baltimore)* 2005;84:377-85.
- Ward MM. Factors predictive of acute renal failure in rhabdomyolysis. *Arch Intern Med* 1988;148:1553-7.
- Veenstra J, Smit WM, Krediet RT, Arisz L. Relationship between elevated creatine phosphokinase and the clinical spectrum of rhabdomyolysis. *Nephrol Dial Transplant* 1994;9:637-41.
- de Meijer AR, Fikkers BG, de Keijzer MH, van Engelen BG, Drenth JP. Serum creatine kinase as predictor of clinical course in rhabdomyolysis: a 5-year intensive care survey. *Intensive Care Med* 2003;29:1121-5.
- Woodrow G, Brownjohn AM, Turney JH. The clinical and biochemical features of acute renal failure due to rhabdomyolysis. *Ren Fail* 1995;17:467-74.
- Knochel JP. Rhabdomyolysis and myoglobinuria. *Annu Rev Med* 1982;33:435-43.
- Zager RA, Gamelin LM. Pathogenetic mechanisms in experimental hemoglobinuric acute renal failure. *Am J Physiol* 1989;256:F446-F455.
- Zager RA. Studies of mechanisms and protective maneuvers in myoglobinuric acute renal injury. *Lab Invest* 1989;60:619-29.
- Zager RA, Foerder CA. Effects of inorganic iron and myoglobin on in vitro proximal tubular lipid peroxidation and cytotoxicity. *J Clin Invest* 1992;89:989-95.
- Reeder BJ, Wilson MT. Hemoglobin and myoglobin associated oxidative stress: from molecular mechanisms to disease states. *Curr Med Chem* 2005;12:2741-51.
- Holt S, Reeder B, Wilson M, et al. Increased lipid peroxidation in patients with rhabdomyolysis. *Lancet* 1999;353:1241.
- Bonventre JV, Weinberg JM. Recent advances in the pathophysiology of ischemic acute renal failure. *J Am Soc Nephrol* 2003;14:2199-210.
- Gabow PA, Kaehny WD, Kelleher SP. The spectrum of rhabdomyolysis. *Medicine (Baltimore)* 1982;61:141-52.
- Hatamizadeh P, Najafi I, Vanholder R, et al. Epidemiologic aspects of the Bam earthquake in Iran: the nephrologic perspective. *Am J Kidney Dis* 2006;47:428-38.
- Sieb JP, Penn AS. Myoglobinuria. In: Engel AG, Franzini-Armstrong C, eds. *Myology*. New York: McGraw-Hill, 2004:1677-92.
- Simerville JA, Maxted WC, Pahira JJ. Urinalysis: a comprehensive review. *Am Fam Physician* 2005;71:1153-62. [Erratum, *Am Fam Physician* 2006;74:1096.]
- Wakabayashi Y, Kikuno T, Ohwada T, Kikawada R. Rapid fall in blood myoglobin in massive rhabdomyolysis and acute renal failure. *Intensive Care Med* 1994;20:109-12.
- Mikkelsen TS, Toft P. Prognostic value, kinetics and effect of CVVHDF on serum of the myoglobin and creatine kinase in critically ill patients with rhabdomyolysis. *Acta Anaesthesiol Scand* 2005;49:859-64.
- Grossman RA, Hamilton RW, Morse BM, Penn AS, Goldberg M. Nontraumatic rhabdomyolysis and acute renal failure. *N Engl J Med* 1974;291:807-11.
- Oh MS. Does serum creatinine rise faster in rhabdomyolysis? *Nephron* 1993;63:255-7.
- Corwin HL, Schreiber MJ, Fang LS. Low fractional excretion of sodium: occurrence with hemoglobinuric- and myoglobinuric-induced acute renal failure. *Arch Intern Med* 1984;144:981-2.
- Better OS, Stein JH. Early management of shock and prophylaxis of acute renal failure in traumatic rhabdomyolysis. *N Engl J Med* 1990;322:825-9.
- Akmal M, Bishop JE, Telfer N, Norman AW, Massry SG. Hypocalcemia and hypercalcemia in patients with rhabdomyolysis with and without acute renal failure. *J Clin Endocrinol Metab* 1986;63:137-42.
- Sever MS, Vanholder R, Lameire N. Management of crush-related injuries after disasters. *N Engl J Med* 2006;354:1052-63.
- Shimazu T, Yoshioka T, Nakata Y, et al. Fluid resuscitation and systemic complications in crush syndrome: 14 Hanshin-Awaji earthquake patients. *J Trauma* 1997;42:641-6.
- Gunal AI, Celiker H, Dogukan A, et al. Early and vigorous fluid resuscitation prevents acute renal failure in the crush victims of catastrophic earthquakes. *J Am Soc Nephrol* 2004;15:1862-7.
- Homsí E, Barreiro MF, Orlando JM, Higa EM. Prophylaxis of acute renal failure in patients with rhabdomyolysis. *Ren Fail* 1997;19:283-8.
- Brown CV, Rhee P, Chan L, Evans K, Demetriades D, Velmahos GC. Preventing renal failure in patients with rhabdomyolysis: do bicarbonate and mannitol make a difference? *J Trauma* 2004;56:1191-6.
- Cho YS, Lim H, Kim SH. Comparison of lactated Ringer's solution and 0.9% saline in the treatment of rhabdomyolysis induced by doxylamine intoxication. *Emerg Med J* 2007;24:276-80.
- Ron D, Taitelman U, Michaelson M, Bar-Joseph G, Bursztein S, Better OS. Prevention of acute renal failure in traumatic rhabdomyolysis. *Arch Intern Med* 1984;144:277-80.
- Richards JR. Rhabdomyolysis and drugs of abuse. *J Emerg Med* 2000;19:51-6.
- Moore KP, Holt SG, Patel RP, et al. A causative role for redox cycling of myoglobin and its inhibition by alkalization

- in the pathogenesis and treatment of rhabdomyolysis-induced renal failure. *J Biol Chem* 1998;273:31731-7.
42. Heyman SN, Greenbaum R, Shina A, Rosen S, Brezis M. Myoglobinuric acute renal failure in the rat: a role for acidosis? *Exp Nephrol* 1997;5:210-6.
43. Ho AM, Karmakar MK, Contardi LH, Ng SS, Hewson JR. Excessive use of normal saline in managing traumatized patients in shock: a preventable contributor to acidosis. *J Trauma* 2001;51:173-7.
44. Zager RA, Foerder C, Bredl C. The influence of mannitol on myoglobinuric acute renal failure: functional, biochemical, and morphological assessments. *J Am Soc Nephrol* 1991;2:848-55.
45. Better OS, Rubinstein I, Winaver JM, Knochel JP. Mannitol therapy revisited (1940-1997). *Kidney Int* 1997;52:886-94.
46. Visweswaran P, Massin EK, Dubose TD Jr. Mannitol-induced acute renal failure. *J Am Soc Nephrol* 1997;8:1028-33.
47. Lameire N, Van Biesen W, Vanholder R. Acute renal failure. *Lancet* 2005;365:417-30.
48. Kellum JA. The use of diuretics and dopamine in acute renal failure: a systematic review of the evidence. *Crit Care* 1997;1:53-9.
49. Evans KJ, Greenberg A. Hyperkalemia: a review. *J Intensive Care Med* 2005;20:272-90.
50. Szpirt WM. Plasmapheresis is not justified in treatment of rhabdomyolysis and acute renal failure. *J Cardiovasc Surg (Torino)* 1997;38:557.
51. Ronco C. Extracorporeal therapies in acute rhabdomyolysis and myoglobin clearance. *Crit Care* 2005;9:141-2.
52. Huerta-Alardin AL, Varon J, Marik PE. Bench-to bedside review: rhabdomyolysis — an overview for clinicians. *Crit Care* 2005;9:158-69.

Copyright © 2009 Massachusetts Medical Society.

CORRECTION

Rhabdomyolysis and Acute Kidney Injury

To the Editor: Bosch and colleagues (July 2 issue)¹ observe that although conventional hemodialysis filters do not remove myoglobin (molecular weight, 17.8 kD), hemodiafiltration with super-high-flux dialyzers may be effective.² We used a hemodialysis prescription with a super-high-flux dialyzer (HCO-1100, Gambro) that efficiently removed molecules of up to 60 kD. In two patients with rhabdomyolysis and acute kidney injury, the mean serum myoglobin clearance with a single dialysis treatment was 59%.³ A 4-hour dialysis treatment cleared myoglobin from the equivalent of 9 liters of extravascular fluid (twice the intravascular volume). The kinetics of myoglobin that we observed were similar to the kinetics of free light chains (25 to 50 kD).⁴

The experience gained in the use of super-high-flux dialysis to remove free light chains in myeloma kidney (or cast nephropathy) should expedite the development of a randomized trial of the removal of myoglobin. A randomized, controlled trial of the use of super-high-flux dialysis to remove free light chains is under way.⁵

Kolitha Basnayake, M.B., B.S.

Paul Cockwell, Ph.D.

Colin A. Hutchison, Ph.D.

University of Birmingham

Birmingham, United Kingdom

k.basnayake@bham.ac.uk

References

1. Bosch X, Poch E, Grau JM. Rhabdomyolysis and acute kidney injury. *N Engl J Med* 2009;361:62-72.
2. Ronco C. Extracorporeal therapies in acute rhabdomyolysis and myoglobin clearance. *Crit Care* 2005;9:141-142.
3. Hutchison CA, Harding S, Basnayake K, Bradwell AR, Cockwell P. Myoglobin removal by high cut-off hemodialysis: in-vivo studies. *J Am Soc Nephrol* 2007;18:250A-250A.
4. Hutchison CA, Cockwell P, Reid S, et al. Efficient removal of immunoglobulin free light chains by hemodialysis for multiple myeloma: in vitro and in vivo studies. *J Am Soc Nephrol* 2007;18:886-895.
5. Hutchison CA, Cook M, Heyne N, et al. European trial of free light chain removal by extended haemodialysis in cast nephropathy (EuLITE): a randomised control trial. *Trials* 2008;9:55-55.

To the Editor: In their review article on rhabdomyolysis and acute kidney injury, Bosch et al. summarize the limited efficacy and prognostic impact of extracorporeal myoglobin removal by standard blood-purification techniques. New options are now possible given the

availability of protein-permeable, high-cutoff filters in Europe (filing for approval by the U.S. Food and Drug Administration is in preparation). These filters are currently under investigation for elimination of nephrotoxic free light chains in cast nephropathy associated with multiple myeloma.¹ With an in vivo molecular cutoff at 45 kD, high-cutoff filters are effective in eliminating the 17-kD molecule myoglobin.² We have used high-cutoff hemodialysis for myoglobin removal in severe rhabdomyolysis. Myoglobin clearance with high-cutoff filters, corrected for membrane-surface area, is up to 20 times as high as myoglobin clearance with standard high-flux hemodialysis in intraindividual comparison. With the use of full-size, high-cutoff filters for 2.1 m² of membrane-surface area (Theralite, Gambro), myoglobin clearances in excess of 70 ml per minute can be obtained, resulting in a rapid and highly effective reduction of the plasma myoglobin concentration.

Nils Heyne, M.D.

Martina Guthoff, M.D.

Katja C. Weisel, M.D.

University of Tübingen

Tübingen, Germany

nils.heyne@med.uni-tuebingen.de

Dr. Heyne reports receiving lecture fees from Gambro. No other potential conflict of interest relevant to this letter was reported.

References

1. Hutchison CA, Cook M, Heyne N, et al. European trial of free light chain removal by extended haemodialysis in cast nephropathy (EuLITE): a randomised control trial. *Trials* 2008;9:55-55.
2. Naka T, Jones D, Baldwin I, et al. Myoglobin clearance by super high-flux hemofiltration in a case of severe rhabdomyolysis: a case report. *Crit Care* 2005;9:R90-R95.

The authors reply: Basnayake and colleagues comment on the potential use of extracorporeal removal of myoglobin with the use of super-high-flux dialysis membranes, and Heyne and colleagues suggest the use of high-cutoff membranes for this purpose. Although the experience with super-high-flux membranes or high-cutoff membranes may be encouraging, it is still limited. A number of questions about myoglobin metabolism, kinetics, and body distribution have not been answered, and these issues may complicate the application of the appropriate extracorporeal treatment in terms of frequency, duration, and intensity.¹ In addition, high-cutoff membranes with reported molecular-weight cutoff values of approximately 50 kD (100 kD in the article by Naka et al.²) may be associated with unwanted losses of albumin or other components that may be dangerous for the patient. We believe that although these techniques are promising, randomized, controlled clinical trials will be necessary before they can be recommended. Therefore, we would emphasize that since conventional

dialysis or standard hemofiltration has not achieved clinically significant myoglobin removal,^{1,3} and the experience with high-cutoff membranes or super-high-flux membranes is limited, at present, these techniques cannot be recommended for the preventive removal of myoglobin in rhabdomyolysis.

In Table 3 of our article, we recommend that volume replacement with normal saline solutions should be used for the prevention or treatment of rhabdomyolysis-induced acute kidney injury. The use of solutions containing bicarbonate is optional because their benefits have not been firmly demonstrated. Although slightly hypertonic bicarbonate solutions have been used by some investigators and are commonly used in some countries as 1/6 M sodium bicarbonate (1.4% sodium bicarbonate),³ we agree with others^{4,5} that they should be isotonic or even slightly hypotonic. Since normal saline, commonly called isotonic saline is in fact slightly hypertonic (154 mmol per liter of sodium and chloride), the alternation with 100 mmol of bicarbonate in 1 liter of 5% dextrose is the most appropriate option if alkalinization is used. If 0.45% saline is to be used, it should be combined with 50 to 70 mmol of bicarbonate (rather than the 100 mmol listed in Table 3 of our article). As recommended in the text, volume repletion and alkalinization in patients with rhabdomyolysis should be monitored by the frequent measurement of levels of urine pH and serum bicarbonate, potassium, and calcium.

Esteban Poch, M.D., Ph.D.

Xavier Bosch, M.D., Ph.D.

Josep M. Grau, M.D., Ph.D.

Hospital Clinic of Barcelona

Barcelona, Spain

jmgrau@clinic.ub.es

References

1. Mikkelsen TS, Toft P. Prognostic value, kinetics and effect of CVVHDF on serum of the myoglobin and creatine kinase in critically ill patients with rhabdomyolysis. *Acta Anaesthesiol Scand* 2005;49:859-864.
2. Naka T, Jones D, Baldwin I, et al. Myoglobin clearance by super high-flux hemofiltration in a case of severe rhabdomyolysis: a case report. *Crit Care* 2005;9:R90-R95.
3. Peltonen S, Ahlström A, Kylävainio V, Honkanen E, Pettilä V. The effect of combining intermittent hemodiafiltration with forced alkaline diuresis on plasma myoglobin in rhabdomyolysis. *Acta Anaesthesiol Scand* 2007;51:553-558.
4. Sever MS, Vanholder R, Lameire N. Management of crush-related injuries after disasters. *N Engl J Med* 2006;354:1052-1063.
5. Vanholder R, Sever MS, Ereke E, Lameire N. Rhabdomyolysis. *J Am Soc Nephrol* 2000;11:1553-1561.